

WHY

Is Land and Water Conservation Important To You & Montana?

Montana is a great place to live, and you can help keep it that way!

ARE YOU RAISING HORSES and wondering why you are having to buy more feed each year as your land's productivity declines, leaving bare ground and weeds?

HAVE YOU HAD THE GOOD FORTUNE to buy a place on a creek and are now frustrated that you aren't permitted to remove the brush so you can see the water? DID YOU JUST FIND OUT that those pretty purple flowers along your fence are noxious weeds and threaten the productivity of your land and your neighbor's land?

As you can see, there's a lot to know about owning and managing land, and you need to know even more if you're raising livestock, too. This booklet will get you started and give you lots of information and ideas for your place. With a little time, a little knowledge and not a whole lot of money, you can have a "picture perfect" place that you can be proud of...and protect Montana's land and water. Remember, we're all part of a neighborhood and our actions can affect others. Refer to the last page for information on how to avoid contamination and infringement on others' rights. The things

Look At What You Have

Any landowner needs a management plan. Before developing your plan - look around, make a sketch, and take a few notes about your property. In your sketch, show or note:

Property boundaries
Fences and corrals

Buildings

Wells (human or stock)

55 Septic system

Streams, wetlands, ponds

Bare ground

Weeds

Lawn, pasture, or crop land

* Trees or shrubs

 Soil type (refer to your county soil survey available from the USDA Soil Conservation Service)

Depth to groundwater (check with well driller)


Neighboring land uses

✓ Flat or sloped ground

The four pastures in this "After" drawing allow better management of livestock grazing and increased forage production. A stockwater tank located in the corral is accessible from all pastures and reduces streambank trampling. Shrub and

tree plantings along the streambank prevent erosion, replace weeds and bare areas, and provide wildlife habitat.

Before You Plan...



Con\$ervation Value\$

that you and your neighbors do can greatly improve the health of our resources...the resources we all appreciate about Montana.

Saves money because your land is more productive over the long term

Ensures better water quality for you, your animals, and your neighbors

Provides wildlife habitat

Produces more grass for grazing


Grows healthier livestock

Improves your property values

Makes your place more attractive

Keeps your neighbors happier

Satisfies your responsibility to care for the land



After You Plan!

What Are Your Property Goals?

What do you want?
What can your land support?

Livestock grazing? How many? Wildlife habitat? Good water quality?

Healthy forest? Native plants? A 4-H project? Something else?

You may find that you have to modify some of your goals because they are not realistic for your property.



A Plan For Your Land

Once you've looked at your property and identified your goals, you need to develop a management plan for reaching your goals. Remember, even if you like things just the way they are, you will need to do something to keep weeds from coming in or to keep the water clean! This booklet provides useful information on developing the many different parts of your management plan.

Contents

Fish?

| Weed Management and Soils | . 2 |
|---|-----|
| Pasture and Irrigation Management | |
| Grazing Management and Livestock Health | 4 |
| Grazing Management and Fencing Options | 5 |
| Streams, Wetlands, and Water Quality Protection | 6 |
| Improving Wildlife Habitat | . 7 |
| Forest Management | . 8 |
| Know Your Responsibilities and Homesite Selection | 9 |



Give Your Land A Health Exam

How much of these do you have on your property?

Healthy ground cover (forest, shrubs, grass, or cropland) Weeds or plants that hold the soil poorly (dandelion, knapweed, cheatgrass)

Bare ground

A lot A little Some A little A lot Some A little

...not by insurance, but by vegetation! Vegetation protects

the soil from erosion by rain,

runoff, and wind. Vegetation

increases water uptake by soils and holds soils in place on slopes and along streams.

You'll need a soil test to find Out. Contact your local Soil

Conservation Service or county

A Soil Profile:

NAME OF STREET

it for testing.

How Fertile Is

Organic

Topsoil

Subsoil

Material

If all of your answers are in the first column, your land earns an "A" for health. If most of your answers are in the second column, it is in average condition. If you have any responses in the third column, your land needs immediate help! Read on to learn about conservation practices that will improve your land's health.

Weed Control Weeds spread fast so regularly look for new weed patches on your property and act immediately to treat them by using one or more of the weed control practices listed below. Team up with neighbors to improve effectiveness. Remember, weed control by itself is not enough. It is also necessary to modify the practices that caused weeds to become established in the first place! Is Your Soil Covered?

Prevention. Good land management will help keep desirable vegetation healthy and weeds under control. Buy only weed-seed-free hay, plant only certified seed, wash your vehicle after being in a weed-infested area, monitor your property, and respond quickly to any new weed infestations.

Biological. Biological control attempts to find something in nature that can weaken or eventually kill a weed plant. Successful bioagents include certain fungi and insects that weaken weeds by attacking seed heads and other plant parts.

Mechanical. Mow weeds annually before they go to seed. Pull small weed patches and weeds near streams by hand.

Livestock Grazing. Graze weeds before they go to seed using sheep, goats, or cattle. Because livestock and wildlife can easily carry and spread weed seed on their coats or in their feces, avoid moving livestock from a weedy area to a weed-free area. Some weed species, if eaten, will make livestock sick.

extension office to find out how to Chemical Herbicides. Herbicides may be expensive, but are effective when applied in the proper amounts and at the proper time of year. Read the label instructions carefully and follow directions. Use chemicals away from water to prevent adverse health effects to you and your animals and to prevent pollution of streams and groundwater. Only licensed users can use restricted herbicides. Call a local farm supply store to find out about hiring custom chemical applicators to spray your weeds. Be sure herbicide will not reach and kill desirable trees and shrubs. Properly dispose of leftover chemicals.

TYPES - Know Your Soil

Soils vary widely, even across your backyard. The type of soil you have will influence:

- What type and how much grass or crops your land can produce
- How quickly water moves through the soil
- If the soil will filter human and animal wastes before they reach groundwater
- How often you need to irrigate
- How much fertilizer is needed
- Possible problems with building foundations
- If the area is a wetland

For information about your soil type, refer to your county's soil survey available from the Soil Conservation Service (SCS) office (listed in your phone book under United States Government, Dept. of Agriculture).

For Help Contact your county weed control district or county extension office to obtain a list of noxious weeds in your local area and recommendations on how best to control them.

► Know Your Weeds **Before They**

- Choke out desirable forage for livestock and wildlife
- Reduce the productivity of your pasture and land
- · Cause water pollution and soil erosion because they're less effective at holding the soil
- Spread RAPIDLY!



Knapweed (Spotted, Russian, and Diffuse)



Leafy Spurge



Whitetop



Dalmation Toadflax



Canada Thistle



Purple Loosestrife



Sulfur Cinquefoil

To distinguish from other native cinquefoils, note that sulfur cinquefoil has many stem leaves, few basal leaves, and long, rightangled hairs along stems.





Is Your Annual Pasture and Hay Production?

| | Fertile Soils | | Poor Soils | |
|--------------|-------------------------|---------------------|-------------------------|---------------------|
| , | FEED (Hay) tons/acre | FORAGE AUMs/acre | FEED (Hay) tons/acre | FORAGE AUMs/acre |
| Irrigated | 2-4 | 3-4 | less than 2 | 1-2 |
| Nonirrigated | 1-2 | 1-2 | .5 or less | .5 |
| Rangeland/wo | odland | .5 | .5 or less | .25 |

These figures are averages and may vary up or down, depending on management.

To Increase Your Pasture Production

A pasture is a grazing area for animals enclosed by a fence. Pastures are often planted to nonnative plant species to increase their production. These pastures may need fertilizing, irrigating, and periodic replanting.

- Develop irrigation (if you have a water right, see page 9). Practice irrigation water management. Under-irrigating will shorten the life of your pasture; over-irrigating wastes energy, water, and your time.
- Fertilize according to SCS and soil test recommendations. Believe the soil test! Overfertilizing is not better and can damage water quality.
- Mow pastures to a uniform 3-inch height after grazing to stimulate equal growth of all plants.
- Drag or harrow to spread nutrient-rich manure.
- · Control weeds.
- Reseed. Contact your local SCS office to determine the most productive seed mixture for your purpose and location.

Consider Custom Farming As A Way To Improve Your Pasture

Many landowners find it too expensive to own their own farm equipment for preparing the soil, seeding, harvesting, or baling. Ask your neighbors if they know of any custom farmers or ranchers in the area who will follow your instructions for improving your pasture.

Q: When do I need to irrigate?

A: Irrigate when the soil moisture drops to about 50 percent of its water-holding capacity in the top 3 feet of soil. Check your soil moisture by squeezing several handfuls of soil taken at 6", 12", and 18" depths in your field. Irrigate before the soil at the 18" depth begins to crumble in your hand, since most of the plants' the roots are above 18".



If there is staining on your fingers from squeezing the soil, wait a couple days and test the soil again. If the soil feels only slightly moist, forms a slightly crumbly ball when squeezed in your hand, and there is no staining, then it is time to irrigate (see picture).

For Help The USDA Soil Conservation Service, an irrigation company, or a consultant can provide assistance in designing an appropriate irrigation system for your property.

► Irrigation Systems

Advantages and Disadvantages


Sprinkler irrigation (includes moveable handlines, moveable wheel line, and center pivot) uses the least amount of water, requires labor to move the irrigation pipe, requires some maintenance, and requires an initial investment. Some operational costs are possible.

Flood irrigation requires lots of water, doesn't spread water evenly across the pasture, requires labor to turn water on or off a pasture, is low maintenance, and is the least expensive (assuming irrigation ditches are already in place).

"Big gun" sprinkler irrigation requires high power costs, applies excess water, and doesn't work well on clay-type soils. Water distribution is only fair and cost is moderate. Minimum labor and some maintenance are required.

► Irrigation Management

Depends on Soil Texture



Loam is a combination of all of these.

► Irrigation How Much and How Often?

| Soil | Moisture to be replaced in the 3 -foot rooting zone | Average Peak Season (July/August) Irrigation Frequency | |
|--------------------------|---|--|--|
| Texture | WHEN SOIL IS AT 50% OF ITS WATER-HOLDING CAPACITY* | | |
| Loamy sand Sandy loam | 1.4" 2.3" | 6 days 9 days | |
| Loam | 3.1" 3.2" | 12 days | |
| Clay loam Clay | 3.1" | 13 days 12 days | |

^{*} These moisture replacement estimates are for an alfalfa/grass hay crop. Amounts may vary for other crops. Irrigation is most important for alfalfa during the seedling stage and immediately after cutting. If your soil depth is less than 3', you'll need to irrigate more often and apply less water.

Q: How long should I irrigate?

A. In general, irrigate sandy soils for short periods (2-3 hours) and clay soils for longer periods (9-12 hours). Ask your farm supply store or local SCS office to recommend the correct size spray nozzle for your soil type and your irrigation system. When it rains, see if the rain has gone deeper than the soil surface before considering it a source of water for your crop.

To determine exactly how long to run your system, first place several pans at various locations under your sprinkler system. Run the system for one hour. Average the depth of the water in the pans. This is your hourly application rate. Next, divide the inches of water to replace by the hourly application rate.

EXAMPLE: Loam needs 3.1" of water replaced in the top 3 feet when it is at 50 percent of its water-holding capacity (see irrigation table). If your irrigation system's application rate is 0.3"/hour, you will need to run your irrigation system for ten and a half hours to deliver 3.1" to the soil, since $3.1" \div 0.3"/hour = 10.5 hours$.

Are Your Grazing Animals **Properly Managed?**

Do you have so little grass in your pastures that your animals consume dirt while trying to graze?

Are your animals browsing on trees, shrubs, fences, or barns?

Are your animals losing weight, or are they overweight?

Do your animals have scruffy coats?

Are your animals prone to colic or respiratory problems?

If you answered "yes" to any of these questions, you need a new grazing program that will provide more grass and healthier animals... and save you money in lower feed costs and lower veterinarian expenses!



Continuous grazing allows weeds to grow where grass roots have been weakened. A less dense leaf canopy allows sunlight to reach invading weeds.

Pasture rotation and good grazing management produces more grass, fewer weeds, and no bare ground.



For a Successful **Grazing Program**

- Eliminate continuous season-long grazing.
- Subdivide large pastures into smaller pastures (see sample grazing designs on next page) and develop a pasture-rotation grazing system.
- Corral livestock and feed them hay until your pasture grasses are 6" to 8" high. Move livestock when 50% of the grass plant has been eaten (3" to 4" height remains). Do not regraze until grasses are at least 6" high (will take 1 to 3 months).
- During winter months, continue your rotation to distribute manure and feed wastes evenly across your pastures or hold animals in a
- Allow long rest periods or use a high-intensity, short-duration grazing system to rejuvenate poor condition pasture.
- Provide a water source for each pasture (see next page).
- Irrigate each pasture (if you have irrigation) immediately after grazing to get plants growing again. Do not graze on wet soils.
- Horses do not need 24-hour access to feed or forage. Their nutrition needs can be met with only a few hours of grazing on good pasture each day. Corral animals for the remainder of the day to prevent overgrazing of plants and extend the forage available in your pastures.
- On a limited acreage, you may have only enough pasture to exercise your animals and will need to feed year-round.

Poor Condition Pastures Cause

- colic and respiratory problems from eating dirt
- weight loss
- parasites
- poor coat

Do You Have Enough Feed and Forage For Your Livestock?

In Montana, livestock are usually grazed May through October during the plants' growing season (if you have enough pasture) and fed hay from November through April.

Forage is what your animals consume by grazing. Forage production is measured in animal unit months (AUMs). One AUM is equivalent to the amount of forage consumed by a 1000-pound

Feed is the hay that you provide an animal when forage is not available. Hay production is measured in tons per acre.

Q. How much feed and forage do your animals need each year? A. Average requirements are listed below, but may vary with season, level of use, and the age and size of the animal.

| | Feed (HAY) Tons/month | FORAGE AUMS OF GRAZING/MONTH |
|---------|-----------------------|---------------------------------|
| 1 cow | .4 | 1.2 |
| 1 horse | .5 | 1.25 |
| 1 sheep | 1 | .2 |
| 1 llama | .15 | .3 |
| 1 goat | .1 | .2 |

- Q. How much feed and forage can your land produce?
- A. See Pasture and Hay Production table on page 3.
- Q. Do your feed and forage requirements balance with your land?

A. To find out, do your own calculations following these examples:

FEED REQUIREMENT: 3 horses $x \cdot \frac{5 \text{ tons}}{\text{month}} \times 6 \text{ months} = 9 \text{ tons hay}$

FEED PRODUCTION: 10 acres (fertile nonirrigated soil) $x \frac{1 \text{ ton}}{\text{acre}} = 10 \text{ tons hay}$

FORAGE REQUIREMENT: 3 horses x $\frac{1.25}{\text{month}}$ x 6 months = 22.5 AUMs

Forage Production: 10 acres (fertile nonirrigated soil) $x \frac{1 \text{ AUM}}{2000} = 10 \text{ AUMs}$


In this example, your land will produce enough hay to feed your animals for 6 months. However, you do not have enough forage (grazing) to meet your animals' needs. To avoid overgrazing your pastures each year:

- Buy additional feed or rent pasture
- Increase your pasture production (see "tips," previous page)
- Improve your grazing management
- Reduce your number of animals
- Seek assistance

How Grazing Affects Root Growth

| | PERCENT GRASS | PERCENT NOOT |
|-------------------------------|---------------|----------------|
| | PLANT REMOVED | GROWTH STOPPED |
| Overgrazing occurs when more | 10% | 0% |
| than 50 percent of the grass | 20% | 0% |
| plant is removed all at once. | 30% | 0% |
| Overgrazing stops root growth | 40% | 0% |
| and reduces grass production. | 50% | 2-4% |
| Look what happens when you | 60% | 50% |
| try to sneak in another 10 | 70% | 78% |
| percent "harvest"50 percent | 80% | 100% |
| of the roots stop growing! | 90% | 100% |
| | | |

Notice how the root mass of these grasses decreases in pastures that range from excellent to







SAMPLE Grazing Schedule For A One Herd Multiple-Pasture System

In Montana, livestock are normally grazed May through October during the plants' growing season. Begin grazing when plants are 6" to 8" in height. Move livestock after 50 percent has been eaten (3"- 4" remains). A minimum of 30 days is needed between grazing periods on irrigated pasture and up to 3 months for nonirrigated pasture. You may need to corral livestock and feed them hay

until the pasture regrows.



Stockwater Development

An Essential Part of Your Grazing and Animal Health Programs

As you divide your acreage into several pastures, establish separate water sources for each pasture or a single water source that is accessible from several pastures.. Clean, fresh water is essential for good animal health. Options for stockwater development include:

- 1. A stock tank or pond (consider how you will keep water from freezing in winter).
- 2. Water gaps on a stream. For small acreages, it is highly recommended that you fence your grazing livestock away from streams to keep manure out of the stream. protect and maintain streamside grasses and shrubs, and control erosion (see Water Quality Protection on page 9).

▶ Sample Designs For A Multiple-Pasture W Grazing System Shelter in corral Water in corral Gate

4

For Help

Obtain publications from county extension offices on livestock production, farming, gardening, and 4-H programs.

Assistance is available from the USDA Soil Conservation Service, conservation districts, and private consultants to:

- Design a grazing system
- Increase hay and pasture production
- Design a livestock waste disposal program
 - Design stockwater developments
 - Help you meet water quality standards



ADVANTAGES

4-STRAND BARBED WIRE

Good control of cattle. Skill and design

for construction readily available.

DISADVANTAGES

Barbed wire may be injurious to horses and llamas. Labor and material costs high. Periodic maintenance required. May be damaged by big game.



WOVEN WIRE

Skill and design for construction readily available. Good control of sheep. Add 2 upper strands of barbed wire for cattle. Labor and material costs high. Some maintenance necessary.



Corral

Pasture

Pasture

fence

W

Choosing The Right Fence


There are many types of fences. Each has

FENCING A Grazing Management

4- to 10-STRAND SMOOTH WIRE

4- to 5-strand good for horses. 8- to 10strand will contain large, exotic animals or keep big game out. Durable.

Labor and material costs high. Periodic maintenance required.




ELECTRIC

JACKLEG


Good for establishing pasture rotation program on small acreages. Lightweight, portable, easy to set up or dismantle before and after irrigation. Less expensive.

Weathers poorly. Don't use in lengths over 1,000 ft. Requires regular maintenance. Needs solar or electric power source.



Aesthetically appealing. Very durable. Withstands heavy snow. Good in areas where it is hard to dig or drive posts. Can be adapted for marshy, wet areas. Low maintenance.


High labor and material costs during construction.



POST AND POLE (RAIL FENCE)

Durable. Withstands heavy snowfall Low maintenance.

High labor and material costs




• Maintenance requirements Weather

HOG PANELS

Can be formed into a small, portable pen. Wheels may be attached to make moving easier. Good for establishing rotation grazing for a couple animals on small

Inexpensive and easy to construct. Appropriate for only a few sheep or other small animals. Should be moved once or twice each day.

Visual impact



Do you have a drainfield or livestock corral less than 100 feet from your drinking well or stream?


Are your streambanks bare of vegetation, eroding, or falling into the stream?

Do your well tests show fecal or nitrate contamination?

If you answered "yes" to any of these questions, you will want to take immediate action to correct the problem. Get help!

Uncertain About the Safety of Your Drinking Water?

The Farm *A* Syst program allows you to assess the potential effects of various farmstead practices on your drinking water supplies. In addition to twelve do-it-yourself worksheets, the program provides suggestions for how you can modify your practices and where to go for help. The quality of your drinking water can affect farm values, as lenders consider the cost of corrective actions or cleanup in sale prices. Contact your county extension agent for more information.



- Establish and maintain shrubs and grasses along streams and around animal confinement areas to trap and absorb pollutionladen runoff before it reaches streams or groundwater.
- Locate corrals and other livestock confinement areas away from streams. Use water gaps or off-stream stockwater tanks to minimize livestock trampling of streambanks.
- Avoid over-irrigation that can cause valuable topsoil, fertilizer, and pesticide runoff.
- Properly dispose of manure, feed, and bedding wastes by spreading on your cropland. Be sure soil is not too wet or frozen to absorb wastes. This will reduce your need for expensive commercial fertilizers.
- Locate corral and septic system downslope of your drinking water well.
- Use farming practices that reduce soil erosion and increase water infiltration, such as: minimum tillage, contour farming, filter strips, and grassed waterways.
- Do not mix, apply, or dispose of weed control chemicals, used motor oil, or other toxic substances near streams or where they can leak into groundwater. Contact your county health department for the best method of disposal in your area.

Riparian Areas

are found along streams, lakes, and wetlands. They are comprised of waterloving plants such as alder, willow, cottonwood, and sedges.

YES



Continuous seasonlong grazing often removes important riparian vegetation and may cause streambank erosion and water quality degradation.

These areas make up less than 5 percent of the landscape, yet contain 75 percent of our plant and animal diversity: turtles, beaver, muskrat, wood duck, songbirds, frogs, insects, aquatic organisms, orchids, lilies, and more. Just about everything you like about these areas depends on leaving them in their natural state.

A Healthy Riparian Area

is the key to a healthy stream system. Lush riparian and wetland vegetation along the water's edge will:

Slow flood flows and reduce erosion and property loss **Secure** food and cover for fish, birds, and other wildlife

Keep water cooler in the summer and prevent ice damage in winter

Reduce water pollution by filtering out sediment, chemicals, and nutrients from runoff

Provide important breeding habitat for birds

Shelter animals during calving, lambing, or fawning

 \pmb{Hold} more water in the soil, slowly releasing it for longer season streamflows and groundwater recharge

Does Your Property Have A Wetland?

Wetlands are protected from land management activities that would destroy them or change their function. Wetlands are determined by specific soil, vegetation, and hydrologic characteristics. Contact the Soil Conservation Service to determine if your wet area is a wetland.

For Help

- The U. S. Fish and Wildlife Service's Private Lands Program funds projects that create, enhance, or restore wetlands (761-5450).
- The Montana Department of Fish, Wildlife and Parks' River Restoration Program funds stream corridor improvements, including fencing and bank stabilization (444-2449).
- County extension offices have lots of water publications, including information on how to test your drinking water quality.
- Request the *Guide to Stream Permitting in Montana* from your conservation district. It lists the laws that must be complied with before initiating any activity in or near a stream, lake, or wetland.
- The Montana Department of Health and Environmental Sciences in Helena will answer questions about state and federal water quality laws (444-2406).

Streams, Wetlands, and Water Quality Protection 6

Is Your Property Attractive to Wildlife?

Are there a variety of vegetation types, such as small grains, tall grasses, shrubs, and trees for food? For cover?

Is there a pond, stream, or stockwater tank available to wildlife? Can wildlife avoid predation from domestic animals, such as cats and dogs?

The more "yes" responses you had, the more likely you will enjoy the company of birds, small mammals, and maybe

even deer and elk.

Upland Game Birds

Provide food. Areas of tall grass, thickets of shrubs, and plots of wheat, barley, and other small grains provide food and habitat diversity for pheasants and other upland birds. When harvesting crops, begin cutting from the center of the field outward to flush birds away. Don't worry about water. These birds get moisture from dew and the food they eat.

Provide nesting areas and cover. Plant tall grass along roadsides and ditchbanks and shrubs along fencelines or as part of a windbreak to provide nesting and cover. Since these birds nest on the ground in the spring, avoid mowing, burning, or using weed control chemicals on your tall grass until birds are out of the nest in mid-June. (Some weeds should be sprayed prior to June 15 to control their spread effectively, so weigh your priorities.)

► Song Birds

Provide food and water. Trees and shrubs can provide seeds, fruits, and berries that birds like. Streams, ponds, or stocktanks can provide water. Place a floating board in stocktanks to prevent birds from drowning while watering.

Provide nesting areas and cover. Song birds require a diversity of vegetation heights (tall grass, shrubs, trees) and a variety of foliage densities (evergreen and deciduous trees) for nesting and safety from predators. Perches of different heights, such as old snags, fences, and telephone poles, are used by many birds (from bluebirds to hawks) for resting and searching for food.

Waterfowl

Provide food. Waterfowl like aquatic plants, small insects, snails, and crustaceans. They also feed on grains and forage.

Provide water. Ponds are a natural for attracting ducks, geese, and other waterfowl. Ponds should have shallow and deep areas and well-vegetated banks. Vegetated islands are the safest and preferred for nesting.

Provide nesting areas and cover. Large 40- to 50-acre areas of tall, dense, undisturbed vegetation near open water are needed for successful nesting. A tangle of dead plants from last year's growth will hide nesting hens from predators. This dense, dead vegetation also creates better temperature and moisture conditions for egg hatching.

Wildlife Habitat = Food+ Water+Cover

Wildlife habitat is being lost as more land is subdivided, bringing houses, people, livestock, dogs, cats, and other intrusions. Landowners can help offset this loss of wildlife habitat by growing a diversity of vegetation that provides food and cover for wildlife.

> FOOD requirements will naturally vary by wildlife species, from the seeds and berries required by birds, to the grasses, forbs, and shrubs preferred by deer and elk.

WATER on or near your property in the form of a pond, stream, or developed stockwater will increase the variety of wildlife you can attract.

COVER is needed for hiding from predators. travel corridors, nesting, and shelter.

for Creating Wildlife Habitat

- Plant a diversity of vegetation types and heights.
- Plant shelterbelts and jence rows with evergreens and fruit-bearing shrubs.
- Leave snags and down, woody material for perching, hiding, and nesting.
- Plant small grains or large-seeded grasses for wildlife food.
- Develop ponds or other watering facilities. If you have too much wildlife of the wrong kind, contact the

Montana Department of Fish, Wildlife and Parks, County Extension, or the U. S. Fish and Wildlife Service for help.

Trout and Other Fish

Provide food and cover. In small streams, the majority of "fish food" comes from the insects and leaves that fall into the stream from overhanging vegetation. Overhanging shrubs, sedges, and grasses also help to keep water temperatures cool in summer and reduce icing in winter.

Provide habitat . Fish need riffles and deep pools to meet all of their food and cover needs at different stages in their lives. The rocks found in riffle areas churn up the water, which adds oxygen and carries insects to the fish hiding behind rocks or under overhanging banks. Deep pools provide the coldest, mostoxygenated water in summer and are least likely to freeze in winter, killing fish.


Deer and Elk

Remember, attracting large wildlife may also mean damage to gardens and ornamental plants. More deer and elk around your home may also attract predators, like mountain lions.

Provide food. Deer and elk are primarily grazers, but also browse on trees and shrubs. Creating openings in the forest will increase grass and shrub growth for big game. In winter, deer and elk look for windblown areas where grasses are exposed—that may be your pasture! After feeding, elk and deer look for thickets of shrubs or stands of trees to rest and stay warm.

Provide cover. When deer and elk feed in the open, they like being no more than 600 feet from trees and brush for hiding. Consider maintaining large areas of dense shrub or trees on your property for hiding and shelter, especially near pastures. Areas of dense timber are cooler in the summer and warmer in winter than open areas. If you want to accommodate deer and elk and need a fence, build a low one with a smooth top wire. This is easier for them to cross.

- For Help To develop a plan for improving wildlife habitat on your property, contact your local USDA Soil Conservation Service office, conservation district office, or visit your library or local bookstores.
- Order trees and shrubs that wildlife prefer from the Montana State Nursery (if you own more than ten acres) or ask your local nursery to suggest some native shrub and tree species adapted for your area.
- Information on pond development is available from the Montana Department of Fish, Wildlife and Parks and the USDA Soil Conservation Service.
- The Montana Department of Fish, Wildlife and Parks has an Upland Game Bird Program that provides limited funding to property owners who want to improve habitat for pheasants and other upland game birds. Program participants must allow some public access for hunting on their property.



Is Your Forest Healthy?

Are your trees free of problem insects. diseases, or animal damage?

Are your trees spaced far enough apart to allow some sunlight to reach the plants growing on the ground?

Is there more than one age or size of tree present (e.g., seedling, pole, mature)?

Is there more than one tree species present?

Do you have scattered, rather than piles of, down woody material?

If you had all "yes" answers, your woodlot is looking good. If not, read on...

Protect Your Home from Wildfire

N

Maintain 30' of green lawn or fire-resistant plants around your home.

Prune the lower branches of trees below 12' to remove "ladder fuels" that can cause a ground fire to become a more destructive and harder-to-control crown fire. Have water and fire-fighting tools available.


Avoid using wood shakes for roofing or storing firewood next to your house.

Contact a USDA Forest Service or Montana Department of State Lands office for publications and videos on protecting homes from wildfire.

► Forest Insects and Disease

PROBLEM

▲ VULNERABLE TREES




GALL RUST OR BLISTER RUSTS


STERN SPRUCE

Grand fir, subalpine fir, species of pine > 6" diameter; spruce or Douglas-fir > 14" diameter


■ Pitch tubes or mass of sap on bark surface or mounds of red-orange boring dust on bark




■ Gall rust forms large swellings on branches and trunks. Blister rust cracks bark open in spring, exposing yellow or orange powdery spores.



 Initially, silky webbing in needles; followed by chewed needles turning brown at tips of branches




■ Witches-brooms form



All sizes and ages of Douglas-fir, grand fir and subalpine fir are most susceptible

Individual trees are dying in the stand, tree crowns thinning; rare east of Continental Divide




Outer bark removed, exposing inner layers grooved with parallel teeth marks

for a Healthy Forest

- Maintain diverse species and ages of trees.
- Reduce losses of trees to problem insects and diseases by removing infected trees and slash as soon as possible.
- Thin trees to improve growth, health, and vigor. Thinning will also increase forage. Leave the largest and healthiest trees.
- Avoid season-long livestock grazing that can compact soils and damage trees from browsing or rubbing.
- Locate access roads away from streams; construct adequate drainage. Seed cut slopes promptly to reduce erosion and water pollution.
- Dispose of heavy accumulations of down woody material to reduce fire hazard. Leave snags (standing dead) and larger downed logs for wildlife and forest nutrient cycling.
- When controlling weeds with chemicals, take special precautions not to kill trees.
- When planting trees, select species adapted to your soil, climate, and particular site. Care for new trees by watering regularly and removing competing vegetation in a 1-foot diameter around trunk. Prevent browsing by livestock and wildlife.


• Seek help when planning a timber sale to get top dollar, handle the various permits needed, and see that the remaining stand is in good shape when the harvest is over





For Help

- The Forest Stewardship Workshop Program will teach you how to do an inventory of your property and develop your own management plan. For workshop details, call Extension Forestry (243-2773).
- The Montana Department of State Lands' Private Forestry Assistance Program gives assistance to western Montana landowners setting up timber harvests and provides information about forestry Best Management Practices (BMP's). Request the Forest Stewardship and Water Quality Guidelines booklet.
- The USDA Soil Conservation Service and local conservation districts can provide assistance in developing a forest land grazing plan.
- Private forestry consultants can conduct forest inventories, set up timber sales, and help you achieve your forest management goals. (A directory of consultants is available from the Montana Department of State Lands.)



PORCUPINE DAMAGE



About to Build?

Is the site in a floodplain or close to a stream? Could your access road cause slumping, scar the hillside, or cause sediment to enter a stream?

Will your prospective homesite disturb wildlife habitat?

Does your neighborhood lack covenants that will protect the land, water resources, and future aesthetics of the area?

If you answered "yes" to all of the questions, WHOA—you have some planning to do.

What Is a "Conservation Easement"?

Montana is a great place to live! As more and more people are visiting, buying land, and moving here, the wide open spaces that make Montana so special are shrinking. You can help keep Montana the 'last best place' by considering a conservation easement on your property.


A conservation easement is a legal document between you and the easement holder that specifies what future uses will and will not be allowed on your property. The easement is attached to the deed for your property and remains with the property forever. The easement holder is responsible for ensuring that the terms of the easement are met in the future. Because some future development options are excluded, property taxes may be less.

If you want to take steps to sustain your land's rural agricultural qualities and to maintain wildlife habitat, contact the Montana Land Reliance, Nature Conservancy, or other local land trust organizations.

Tips For Planning A Homesite

- Plan for minimum impact before building.
- Site homes and roads away from streams, on stable soils, and avoid steep slopes.
- · Avoid disturbing wildlife corridors, wetlands, and riparian areas.
- Control your pets so they don't disturb or attract wildlife.
- Maintain or plant native vegetation.
- As a neighborhood working together, you can provide the vegetation diversity that birds, butterflies, and small and large mammals need for food,

cover, and nesting: (a) plant small corner wood-lots, (b) establish shelterbelts edged with shrubs along property boundaries, (c) connect with meadows of native grasses or pasture land, (d) locate house and lawn in a corner of your property to minimize wildlife disturbance.



WHAT You Need To Know As A Montana Landowner

WATER RIGHTS - You must have a water use permit before diverting, withdrawing, impounding or distributing any surface water (or groundwater at rates of 35 gallons per minute or more).

PROTECTION OF STREAMBED AND BANKS - You must have a permit before doing any activity that modifies the stream channel or streambanks.

FLOODPLAIN PROTECTION - You must have a permit before doing any construction work in an area that would be inundated in a 100-year flood. Find out if you are in a floodplain.

CONTROL OF NOXIOUS WEEDS - All counties have laws requiring you to control noxious weeds. Find out which weeds are noxious in your county and how best to control them.

SEPTIC SYSTEM INSTALLATION - Counties regulate septic system installation, including the minimum acceptable distance between your septic system and drinking wells, streams, and groundwater. Counties also approve the septic system design, capacity, and type of soil used to

CITY/COUNTY ZONING - Before building, contact your city (if you are within city limits) or your county planning office to obtain a zoning compliance permit.

WATER QUALITY PROTECTION - You are responsible for preventing livestock manure, pesticides, sediment and other pollutants from reaching waterways.

WETLANDS PROTECTION - You must have a permit to fill, drain, or dredge any waters of the U.S., including wetlands.

STOCKING FISH IN YOUR POND - You will need a permit to stock any species of fish in a

FOREST PRACTICES - In streamside areas, seven forest practices (clearcutting, burning, road construction, etc.) are prohibited on timber sales within a zone of 50 to 100 feet on each side of a stream, lake, or other body of water. A hazard reduction agreement is required before harvesting

OTHER LAND AND WATER PROTECTION MEASURES - Find out if your city or county has special ordinances, such as a Sediment and Erosion Control Ordinance that limits erosion resulting from construction, timber harvest, farming, etc., or an Aquifer Protection Ordinance that protects drinking water wells from contamination, or any type of ordinance that may affect your proposed

AIR QUALITY PROTECTION - Counties determine the best time of year for open burning to minimize deterioration of local air quality and may restrict the use of woodstoves & fireplaces.

OPEN RANGE - Montana is an open range state. Adjacent landowners are equally responsible to maintain the fences between them. As a general rule, fence your property to keep range livestock out and your animals in. It is unlawful for dogs and other predators to harass, kill, or wound cattle, horses, sheep and other livestock.

WHO To Contact

· Water Rights Bureau of the Department of Natural Resources and Conservation, Helena

444-6610

County Conservation District

· U.S. Army Corps of Engineers, Helena

444-6670

· County Floodplain Coordinator or the Floodplain Mgmt, Section. Dept. of Natural Resources and Conservation 444-6654

County weed control district

· County extension office

· County health department or planning office

· City or county planning office

· Water Quality Bureau of the Department of Health and Environmental Sciences (DHES), Helena

444-2406

· U.S. Army Corps of Engineers, Helena

444-6670

· Water Quality Bureau/DHES, Helena

444-2406

 USDA Soil Conservation Service Field Offices (to identify wetlands) · Dept. of Fish, Wildlife and Parks, Helena

444-2449

· Dept. of State Lands, Missoula

542-4300

· City/county planning office

- · Water quality district
- Conservation district
- · County health department or local fire department
- · Department of Livestock, Helena

444-2023

Inquire about other laws that may apply to your property or proposed activities

Know Your Responsibilities & Homesite Selection



- Productive agricultural landWildlife
 - Clear streams
 - Native plants
 - Healthy forests

About This Publication

PROJECT COORDINATOR:

Joan Schumaker, Resource Specialist, Montana Department of Natural Resources and Conservation

DEVELOPED IN COOPERATION WITH:

Missoula County Conservation District Montana Riparian Wetland Association

Montana State University Extension Service USDA Soil Conservation Service*

WITH SPECIAL ASSISTANCE FROM:

Sady Babcock and Tara Comfort, Soil Conservationists, Missoula County Conservation District

Meg Bishop, Dan Himsworth, Amy Smith, Sandy Smith, Tim Wiersum, and Wendy Williams, USDA Soil Conservation Service

EDITED BY:

Carole Massman, Publications Manager, Montana Department of Natural Resources and Conservation

COOPERATIVE FUNDING PROVIDED BY:

Bitterroot Conservation Distric

Departments of Plant, Soil and Environmental Sciences and Political Science, Montana State University

Flathead Conservation District

Gallatin Conservation District

Greater Yellowstone Coalition

Lake County Conservation District

Lewis and Clark County Conservation District

Missoula County Conservation District

Montana Department of Health and

Environmental Sciences

Montana Department of Natural Resources

and Conservation

Montana Department of State Lands

Montana Fish, Wildlife and Parks

Montana Riparian Wetland Association

Education Committee

Northwest Area Foundation

US Environmental Protection Agency

USDI Fish and Wildlife Service.

Montana "Partners for Wildlife" Program

USDA Forest Service Northern Region

USDA Soil Conservation Service

Yellowstone Conservation District

IDED BY: PHOTO CREDITS:

Dr. Susan Hagle, Scott Tunnock, Ken Gibson, and Steve Chadde, USFS Northern Region
Kitty Knaphers, Cascade County Weed District
Marcia Rueter Leritz and Terry Lonner, Media Works
Montana Forest Stewardship Program
Montana State University Publications
Peter Rice, University of Montana
Jim Schoenbaum, Missoula County
Wyoming Weed and Pest Control

DESIGNED AND ILLUSTRATED BY:

Marcia Rueter Leritz, Graphic Artist Martha A. Lonner, Production Manager Media Works, Bozeman, Montana

PRINTED BY:

Color World, Bozeman, Montana

To Request Copies:

Conservation Districts Bureau
Department of Natural Resources
and Conservation
P. O. Box 202301
Helena, MT 59620-2301
(406) 444-6667

* NOTE: The name of the USDA Soil Conservation Service (used throughout the booklet) has since been changed to the USDA Natural Resources Conservation Service.



You may reproduce or copy any portion of this booklet by notifying the Montana Department of Natural Resources and Conservation. Please acknowledge this publication as the source.